

Innovations in Nano Bioenergy
and Bioprocess Technology

Series Editor :Neha Srivastava

BIOFUELS PRODUCTION USING SUSTAINABLE BIOPROCESSING TECHNOLOGIES

VOL 4



EDITORS

DAN BAHADUR PAL and ASHISH KAPOOR

Biofuels Production Using Sustainable Bioprocessing Technologies



NIPA® GENX ELECTRONIC RESOURCES & SOLUTIONS P. LTD.

New Delhi-110 034

About the Series Editor



Dr. Neha Srivastava is working as Research Scientist at Department of Chemical Engineering and Technology, IIT (BHU) Varanasi, India with 14 years research experience. She is globally recognized researcher in the area of waste biorefinaries, bioprocess & microbial technology and green nanotechnology. She has published over 100 SCI and SCOPUS indexed publications and currently serving as “Editor In Chief” of the International Journal Sustainable Process Connect. She is editor of the Nature Journal Scientific Report and Frontiers in Microbiology. She has published 28 books with current running 08 book series as “Series Editor” and has 3 patents in her credit. Dr. Srivastava also ranked world’s top 2% scientist by Stanford University’s study since 2021 with current H-index 40 and 5500 citations.

Scan for Other Volumes of the Series



About the Volume Editors

Dr. Dan Bahadur Pal is currently working as an Assistant Professor, Department of Chemical Engineering, Harcourt Butler Technical University, Kanpur-208002, Uttar Pradesh India. He has received his M. Tech and Ph.D. in the field of Chemical Engineering from Indian Institute of Technology (BHU) Varanasi, Uttar Pradesh, India. Before that, he has completed his B. Tech in Chemical Engineering from UPTU, Lucknow. Dr. Pal's research interest is nano-technology, catalysis, energy and environment and waste management with a special focus in developing process and materials by using waste as raw materials. He also prefers to work on bio-waste processing and value addition. Dr. Pal published more than 92 publications in reputed journals and books along with 40 book chapters and 9 book editing.

Dr. Ashish Kapoor has received his B. Tech and M. Tech degrees in Chemical Engineering from Indian Institute of Technology Madras, Tamil Nadu, India. He received his doctorate in Chemical Engineering from University of Illinois at Urbana-Champaign, USA. His research interests include environmental remediation, microfluidic sensors for water quality monitoring, transport processes, modeling and simulation. He has published more than 45 publications in reputed journals and 10 book chapters. He has been granted one patent.

Innovations in Nano Bioenergy and Bioprocess Technology Volume 4

Biofuels Production Using Sustainable Bioprocessing Technologies

Series Editor

Neha Srivastava

Research Scientist

Department of Chemical Engineering and Technology
IIT (BHU) Varanasi, Uttar Pradesh, India

Volume Editors

Dan Bahadur Pal

Department of Chemical Engineering
Harcourt Butler Technical University
Kanpur, Uttar Pradesh 208002, India

Ashish Kapoor

Department of Chemical Engineering
Harcourt Butler Technical University
Kanpur, Uttar Pradesh 208002, India



NIPA® GENX ELECTRONIC RESOURCES & SOLUTIONS P. LTD.
New Delhi-110 034



**NIPA® GENX ELECTRONIC
RESOURCES & SOLUTIONS P. LTD.**

101,103, Vikas Surya Plaza, CU Block
L.S.C.Market, Pitam Pura, New Delhi-110 034
Ph : +91 11 4386 0225, 9717133558, 9540816132
E-mail: newindiapublishingagency@gmail.com
Website: www.niparesources.com

© 2025, Publisher

Print ISBN: 978-93-7219-053-3

ebook ISBN: 978-93-7219-642-9

All rights reserved. No part of this publication may be reproduced, stored in a retrieval system or transmitted in any form or by any means, including electronic, mechanical, photocopying recording or otherwise without the prior written permission of the publisher or the copyright holder.

This book contains information obtained from authentic and highly reliable sources. Reasonable efforts have been made to publish reliable data and information, but the author/s, editor/s and publisher cannot assume responsibility for the validity, accuracy or completeness of all materials or information published herein or the consequences of their use. The work is published with the understanding that the publisher and author/s are not attempting to render any professional services. The author/s, editor/s and publisher have attempted to trace and acknowledge the copyright holders of all material reproduced in this publication and apologize to copyright holders if permission and/or acknowledgements to publish in this form have not been taken. If any copyrighted material has not been acknowledged, please write to us and let us know so that we may rectify the error, in subsequent reprints.

Trademark Notice: NIPA®, the NIPA® logos and their presentations (the way they are written/presented) in this book are the trademarks of the publisher and hence may not be used without written permission, if copied or used without authorization, the infringer will be prosecuted as per law.

NIPA® also publishes books in a variety of electronic formats. Some content that appears in print may not be available in electronic books, and vice versa

Composed and Designed by NIPA®.

Preface

This book provides a comprehensive overview of bioprocess technologies designed to transform renewable resources into biofuels. A major focus is placed on waste valorization, specifically the conversion of agricultural residues and food waste into biofuels like bioethanol, biobutanol, and biohydrogen. By tapping into waste streams, the book emphasizes the importance of a circular economy and the role of resource recovery in sustainable bioenergy production. It highlights the potential of algae as a biofuel source, while addressing the challenges and prospects for large-scale production. The text explores the role of machine learning and genetic engineering, showcasing how these technologies optimize biofuel processes and drive innovation in the field. The book also discusses advanced energy management systems, aiming to enhance the sustainability of biofuel production through energy recovery and integration. Microbial biofuel production and enzyme-based processes are examined in detail, offering insights into cutting-edge techniques that harness biological activity for bioenergy generation. In addition to these practical approaches, the text delves into the use of computational Modeling for optimizing biofuel production. The focus on integrated biorefineries highlights their role in creating efficient, scalable systems, which are essential for a sustainable bioeconomy.

In Chapter 1, we start with an Algal Biofuel Utilization: Challenges and Prospects. Chapter 2 looks at how Production of Bioenergy using Algal Biomasses: Challenges and Prospects. Chapter 3 focuses on how the Exploring the Biofuel Production Through Algal Genetic Engineering: Current Techniques and Challenges, while Chapter 4 highlights the role of Smart Energy Management and Recovery Towards Sustainable Bioenergy, Chapter 5 introduces the Pre-Treatment of Lignocellulosic Wastes for Biofuel Production. Chapter 6 Valorization of agricultural wastes for Glycerol production. Chapter 7 explains how Valorization of Agricultural Wastes for Bioethanol Production, Chapter 8 discusses how Smart fuel and its Management Towards Sustainable Bioenergy, Chapter 9 looks at how Cellulosic Biomass Fermentation for Bioenergy Generation, Chapter 10 explores how Lignocellulosic biomass biorefineries towards circular economy.

Chapter 11 explains how Advances in Green Microbial Biofuel Production and Chapter 12 focuses on Lignocellulosic Biomass for the Production Biohydrogen.

This book aims to simplify complex ideas and provide practical insights into how biomass can revolutionize biofuels, helping to meet the world's growing energy needs in an environmentally friendly way.

Editors

Acknowledgement

We sincerely thank all contributors and researchers whose expertise enriched this book. The editors are grateful to the Harcourt Butler Technical University, Kanpur, India for technical facilities. We are grateful to the publishers for their support and to the academic community for providing the foundation of knowledge in this field.

Special thanks to our families, friends, and colleagues for their encouragement, and to our readers, whose curiosity inspires such efforts. We hope this book serves as a valuable resource for further exploration in Biofuels production using Sustainable bioprocessing.

Editors

Contents

<i>Preface</i>	v
<i>Acknowledgement</i>	vii
1. Algal Biofuel Utilization: Challenges and Prospects	1
<i>Priyanka Yadav, Diksha Singh, Shivani Mishra, Pratibha Tiwari Mayank Saxena, Ravi Yadav and Kusum Yadav</i>	
2. Production of Bioenergy Using Algal Biomasses: Challenges and Prospects	19
<i>S. Shemaiah, S. Vishali, E. Poonguzhali and S. Kiruthika</i>	
3 Exploring the Biofuel Production Through Algal Genetic Engineering: Current Techniques and Challenges	43
<i>Anand Kumar and Ankit Agrawal</i>	
4. Smart Energy Management and Recovery Towards Sustainable Bioenergy	65
<i>Preeti Chaurasia, Uma Chaurasia, Ashutosh Kumar, Deepak Kumar and Aniruddh Kumar</i>	
5. Pre-Treatment of Lignocellulosic Wastes for Biofuel Production	87
<i>Dilip Kumar, Parichay Gupta, Shruti Gupta, Manish Singh Rajput and Mahendra Kumar</i>	
6. Valorization of Agricultural Waste for Sustainable Glycerol Production: Technological Pathways, Challenges, and Future Prospects	107
<i>Kavitha E, Vishali Solaippan and Sam David Swaminathan</i>	
7. Valorization of Agricultural Wastes for Bioethanol Production ..	133
<i>S. Shemaiah, S. Kiruthika, S. Sam David and E. Poonguzhali</i>	
8. Smart fuel and its Management Towards Sustainable Bioenergy	157
<i>S. Vishali, S. Sam David and E. Kavitha</i>	

9. Cellulosic Biomass Fermentation for Bioenergy Generation	179
<i>S. Kiruthika, Sruthi J Nair, E. Poonguzhali and Kavitha Edward</i>	
10. Lignocellulosic Biomass Biorefineries Towards Circular Economy	205
<i>Meenu Mariam Jacob, Muthamilselvi Ponnuchamy, Dan Bahadur Pal and Ashish Kapoor</i>	
11. Advances in Green Microbial Biofuel Production.....	225
<i>Reena Saxena, Shreyasha Mishra, Rajeev Kumar Upadhyay and Rahul</i>	
12. Lignocellulosic Biomass for Biohydrogen Production.....	245
<i>Vishali Solaiappan, Shemaiah Sam, Kiruthika Sathiasivan and Kavitha Edward</i>	
<i>Index.....</i>	265

Exploring the Biofuel Production Through Algal Genetic Engineering: Current Techniques and Challenges

Anand Kumar and Ankit Agrawal

Department of Biotechnology, IFTM University, Moradabad, Uttar Pradesh

Abstract

In this era, the consumption for fuel has surged, leading to the exploitation of natural resources and increased pollution. This condition has magnified the need to discover safe alternative energy sources. Algae are known for their significant role to produce a broader range of valuable products, including biomolecules like lipids and carbohydrates that can be transformed into biofuels. This has captured more attention to their use. Algae are considered as more economic biofuel rather than traditional crop-based sources which consume arable land and harm food security. But due to their limited capability of manufacturing the biomolecules constraints the production of biofuel. To overcome this situation, numerous genetic engineering techniques have been developed to enhance yield of biofuel. Recent innovations like CRISPR/Cas9, RNA interference (RNAi), transgenic methods, genome sequencing, and micro biome engineering, activates more genetic modifications in algae. Additionally, to support the fourth-generation biofuels, genetic engineering and computational methodology are essential for standardizing algal growth conditions. This chapter provides insights to enhance biofuel production from algae, by creating the recombinant microalgal strains through genetic engineering methods. To overcome preexisting production hurdles, techniques like omics and flux analysis facilitate the mathematical evaluation of such modifications. Apart from to improve biomass and lipid content in certain algal species, technologies such as: high-throughput strain engineering, omics-informed modeling and microbiome engineering are playing key role. This collaborative use of technologies opens the gateway for comprehensive strategies in efficient algal biofuel production.

Keywords: Microalgae, Genetic modification, Biofuels, RNAi, Computational biology and Renewable energy

1. Background

The growing global population, industrialization, and urbanization have driven rising energy demand, with primary energy use increasing by 5.8% in 2024 (BP Statistical Review of World Energy 2024). The need for secure, affordable, and low-carbon energy is critical due to fossil fuel depletion, global warming, and geopolitical tensions. Clean energy sources like solar, wind, and biofuels are being explored, with biodiesel standing out as the only biofuel capable of fully replacing fossil fuels (Aizouq et al., 2020). Many countries, including India and China, are advancing biofuel production, with the latter viewing it as a key to energy security. Biofuels, derived from various sources like algae, crops, and waste, include biodiesel, bioethanol, and biogas. They are categorized into four generations based on feedstock. First-generation biofuels, made from edible materials, face food vs. fuel competition. Second-generation biofuels from non-edible sources are more sustainable but still face resource limitations (Lv et al., 2019). Third- and fourth-generation biofuels from microalgae offer higher yields and better land use, with species like *Chlorella vulgaris* and *Nannochloropsis oceanica* showing promise for biodiesel production. Microalgae's fast growth and efficiency have led to advances in biodiesel production, although challenges remain, such as species selection and production scale-up (Arora et al., 2021). But there is a constraint for commercial availability of these biofuel which are the economic and environmental obstacles that are creating problematic condition for approval and use of genetically engineered microalgae for their potential to enhance lipid yields. This chapter delves into the current opportunities and challenges in microalgae biofuel production, focusing the need for regular research and innovation (Maliha and Abu-Hijleh 2022).

2. Choosing the Different Classes Of Microalgal Strains

The achievement of huge-scale microalgal culture relies on picking appropriate species. The production of various lipids greatly affects the selection of strains and biofuel properties. For the biodiesel, Triacylglycerol (TAG) is the preferred lipid but compared to green algae its content is typically low in cyanobacteria. with the average lipid content of 25.5%, green algae (Chlorophyceae) stand as the largest class of oleaginous microalgae, which can rise under nutrient stress (Saini et al., 2021). Also, the biomass productivity is equally important for high lipid productivity as high lipid content is beneficial. For biodiesel production, strain selection should focus on both lipid content and biomass. *Chromochloris zofingiensis* shows the highest lipid productivity among all classes of green algae. The saturated (SFA) and monounsaturated fatty acids (MUFA) fatty acid (FA) being dominant, their composition also plays a vital role in biodiesel quality. It has seen those high levels of C16:0 and C18:1 in microalgae is ideal for biodiesel production (Bharti et al., 2021). Therefore, for improved biodiesel performance, strain selection should consider both lipid productivity and FA composition (Bagchi and Mallick, 2016).

3. Microalgae Cultivation Systems Stimulates Lipid Accumulation

As the microalgae cultivation can use basic nutrients or wastewater while absorbing CO_2 it is relatively simple because it can occur in open or closed systems. It is interesting that systems like artificial ponds and large water bodies are used in open systems with various agitation methods, while photobioreactors (PBRs) like plastic bags, tubes, and plate reactors are used in closed systems (Plohn et al., 2021). Due to lack of light inefficiency, CO_2 loss stable environmental control and easy contamination, initially open systems were preferred commercially for lower costs and ease of operation (Behl et al., 2020). These issues are addressed properly in closed PBRs, offering reduced contamination, better photosynthetic efficiency and higher biomass and lipid productivity (Wang et al., 2021).




Fig. 1: Graphical representation of time laps for growth of biofuels

3.1 Factors Affecting Lipid Accumulation

Oleaginous microalgae are abundant, but lipid levels vary based on lipid metabolism modifications. Common methods to increase lipid productivity include nutrient deficiencies, stress, and environmental changes (Hui et al., 2016). Phototrophic microalgae produce biomass by absorbing CO_2 , with 1–5% CO_2 concentration being optimal for growth. Higher CO_2 concentrations inhibit growth, while low or excessively high concentrations reduce lipid content (Nakanishi et al., 2014). In this range, increased CO_2 helps species like *Tribonema minus*, *Chlamydomonas*, and *Nannochloropsis* accumulate more lipids. For large-scale production, industrial flue gas provides an economical CO_2 source for biofuel production (Krishnan et al., 2014). Microalgae species thriving in elevated CO_2 environments, with some showing enhanced lipid accumulation at 15% CO_2 , have been identified. Chile and Colombia have implemented microalgae photobioreactors for carbon capture, demonstrating the feasibility of using high- CO_2 -adaptive microalgae for large-scale biofuel production (Yanlin et al., 2014). However, this represents only a limited selection of species. NaHCO_3 is a valuable, easier-to-handle carbon source for microalgae compared to gaseous CO_2 (Bagchi

and Mallick, 2016). Studies show that NaHCO_3 significantly increases lipid content in species like *D. salina*, *D. tertiolecta*, *Scenedesmus*, and *Nannochloris*, especially under nutrient-deficient conditions (Shin et al., 2018). Some microalgae even prefer NaHCO_3 over CO_2 for lipid biosynthesis, likely due to enhanced enzyme activity in lipid metabolism, which boosts inorganic carbon assimilation.

3.2 Performance of Photobioreactors

PBRs allow for a broader range of microalgae species compared to open systems, enabling higher biomass and lipid accumulation. Species like *C. reinhardtii*, *N. oceanica*, and *Chlorella pyrenoidosa* thrive in PBRs, showing better growth performance than in open systems (Aizouq et al., 2020). The design of PBRs significantly impacts productivity, leading to innovations to optimize biomass production, reduce costs, and improve efficiency, especially through membrane PBRs using wastewater for lipid production. PBRs enhance light energy efficiency through improved light paths and mass transfer, boosting biomass and lipid accumulation. Hydrodynamics and fluid mixing in PBRs are critical for nutrient delivery and preventing sedimentation, with optimized designs improving biofuel production (Arora and Philippidis 2021). Proper reactor configurations and operating conditions are crucial to maximizing productivity while preventing cell damage (Bergthorsen and Thomson, 2015)

Limitations and Future Perspectives of PBRs

Although various PBR configurations are effective for microalgae growth, they are often energy-intensive and costly. For example, flat plate PBRs require large space and are prone to photoinhibition and high costs (Bharti et al., 2021). Column PBRs have lower surface-to-volume ratios and are costly to mix, while tubular PBRs face overheating and fouling issues (Chong et al., 2023). Softframe PBRs struggle with fragile materials, poor mixing, and leakage. Hybrid PBRs, despite their potential, encounter membrane fouling and energy imbalance. Scaling up from pilot plants to full-scale operations is difficult without consistent data on both biomass and lipid productivity. Optimizing PBR design for different microalgae species and metabolites is essential for future sustainable development and scaling (Behl et al., 2020).

4. Opportunities in Converting Microalgae to Biofuel and Its Scale-up

Transportation fuels and the energy industry contribute most energy-related emissions. Renewable energy currently makes up about 11% of global primary energy but is expected to reach 60% by 2070. Early biofuel development focused on converting plant sugars into ethanol and upgrading vegetable oils (Noraini et al., 2014). While biofuel production and use are expected to increase, its contribution to total energy demand, especially in transportation, will remain limited due to competition with food and fiber production, agricultural practices in emerging markets, market constraints, and biodiversity conservation. Research now centers on finding sustainable biofuel feedstocks, scaling up microalgae use, and utilizing

hydrothermal liquefaction, though challenges exist in upscaling algae cultivation, particularly in marine and coastal environments (Bergthorsen and Thomson, 2015). The technology to make each stage of biofuel production from microalgae economically viable—such as cultivation, harvesting, transport, pretreatment, and conversion—has not yet been developed. Key challenges include securing sufficient fresh water, nutrient sources (nitrogen, phosphorus, trace elements), and cost-effective, energy-efficient methods for biomass harvesting, oil extraction, and conversion. Additionally, technologies for CO₂ mitigation and system integration are still needed. If these issues are resolved, algal cultivation could provide a sustainable feedstock and offer potential for CO₂ remediation, with microalgae fixing more CO₂ than terrestrial biomass (Ullah et al., 2014). During the conversion process, organic nitrogen turns into ammonia in reducing environments and NO_x in oxidizing environments. In biogas production, high nitrogen content in algal biomass can cause ammonia toxicity, hindering bacterial decomposition (Coelho et al., 2014). Nitrogen also produces NO_x during gasification, which, when performed with limited oxygen, releases NO_x into the atmosphere—a concern due to its greenhouse gas properties, requiring strict emission controls. Challenges in scaling up the microalgae industry include high installation and operating costs, strain robustness, lipid quality for biodiesel, lipid loss during scale-up, and difficulties in managing outdoor cultivation conditions (Garcia-Moscosa et al., 2013).

5. Biofuels Produced from Algae

5.1 Biohydrogen

Microalgae have been used as feedstock for producing biodiesel, bioethanol and biogas. Various microbial species have been used as feedstock for biohydrogen production, among which *Chlorella* sp., *Scenedesmus* sp. and *Saccharina* sp. have been extensively studied (Bilad et al., 2021). To enhance the hydrogen production efficiency, different pretreatment methods were explored (Wieczorek et al., 2014).

5.1.1 Hydrogen Production from Physically and Chemically Pretreated Microalgae

Physical and chemical pretreatments such as mechanical, heat, ultrasonic, acid, base, and ozonation are commonly used to disrupt microalgal cell walls and enhance subsequent biological conversion processes (Bra'nyikova et al., 2011). For example, Ortigueira et al. investigated hydrogen production from dry ground *Scenedesmus obliquus* biomass. The effectiveness of microalgae as a fermentable feedstock depends on the recovery of intracellular sugars and those in the cell walls. Thermal pretreatment often involves additional processes, like pressure build-up with autoclaving or electromagnetic radiation with microwaving (Cheng et al., 2014). Chemical pretreatments have also been shown to improve hydrogen production. Physically and chemically treated microalgae yield higher hydrogen compared to untreated algae, with the highest yields (94.3–338 mL H₂/g VS) seen in heat-treated algae, particularly *Chlorella sorokiniana* (Ortigueira et al., 2015).

5.1.2 Hydrogen Production From Biologically Pretreated Microalgae

Biological approaches use microbes and enzymes to disrupt biomass and release intracellular materials, boosting biohydrogen production (Chong et al., 2023). Enzyme selection depends on the cell wall composition, and factors like pH, temperature, and the enzyme/microalgae ratio are critical for optimizing enzymatic treatment. Inappropriate pH can denature enzymes, while high temperatures may increase enzyme-microalgae interactions until denaturation occurs. A high microalgae-to-enzyme ratio can increase viscosity, hindering enzyme activity. Microalgae cell walls contain cellulose, mucopolysaccharides, and peptidoglycans, making macerozyme, cellulases, and lysozymes useful for disruption (Chong et al., 2021). Enzyme-treated microalgae produced higher hydrogen yields (11–135 mL H₂/g VS) compared to microbial consortium-treated algae, and combining different enzymes significantly improved hydrogen yield. Combined pretreatment methods, typically involving a physical and a chemical treatment, are used to efficiently disintegrate microalgal biomass and enhance hydrogen production (Coelho et al., 2014). The most common combination is heat and acid pretreatment, but other combinations, such as heat with base, enzymatic treatment, oxidizing agents, or ozone with ultrasonication, have also been applied. Some methods even use three or more treatments, like acid-heat-enzyme or acid-microwave-enzyme pretreatment. These combinations improve hydrogen yield, which ranges from 33.56 to 958 mL H₂/g VS. Acid-heat pretreatment shows the highest potential, with *Chlorella* species yielding the most hydrogen (Roy et al., 2014).

5.2 Starch Derived Alcohols

Ethanol production from algae relies on fermenting algal polysaccharides such as starch, sugar, and cellulose. Microalgae can contain up to 70% carbohydrate content, mainly starch, under specific conditions. Their cell walls consist of an inner layer with cellulose and hemicellulose, and an outer layer containing polysaccharides like pectin, agar, and alginate, though composition varies by species (Dubini et al., 2009). Both microalgal starch and cell wall polysaccharides are fermentable for bioethanol production. Macroalgae also contain significant carbohydrates (25-60%), with species like *Ascophyllum*, *Porphyra*, and *Palmaria* having the highest polysaccharide content. Their cell walls mainly consist of cellulose and hemicellulose, comprising 2-10% of dry weight. Lignin is found only in *Ulva* species, constituting 3% of dry weight (Wang et al., 2010). Unlike microalgae, macroalgae contain abundant polymers like alginate, mannitol, glucan, and laminarin. Alginates, extracted from brown algae, are used in pharmaceuticals, paper, adhesives, oil, photography, and textiles. Carrageenan and agar, derived from red algae, are used as stabilizers and gelling agents, primarily in the food industry (Fawzy et al., 2021). Mannitol, found in brown algae, varies with season and environment, and is used in pharmaceuticals, paint, leather, and food industries. Laminarin boosts immunity and offers protection against infections and radiation. Ulvan, found in *Ulva*, is a sugar source for fine chemical production (Kumar et al., 2016).

5.3 Diesel Fuel Surrogates

Biodiesel feedstocks, such as vegetable oils, animal fats, and microalgal oil, are more viscous than petro-diesel and require conversion to meet regulatory standards. Conversion methods include direct use, blending with petro-diesel, micro-emulsions with solvents or alcohols, pyrolysis, and transesterification (Fuad-Hossain et al., 2020). Transesterification, the most common method, has been used since before World War II. However, it is energy-intensive, increases production costs, and involves challenges like catalyst removal, saponification, and glycerol separation, which complicate the process and reduce yield. Enzymatic esterification is an effective method for converting oils with high free fatty acid content to alkyl esters without solvents, offering benefits like lower energy consumption and easier product recovery (Garcia-Moscosa et al., 2013). Yet, due to issues with enzyme production rates and leftover reactions, hurdles remain in making the process efficient for large-scale biodiesel production. To achieve powerful alternative, offering sustainable, renewable, and ecofriendly biofuels high-tech thermal conversion processes have raised with economic potential. But the big-scale production is concise because there is less significant testing and technology transfer methods (Cheng et al., 2014).

5.4 Alkanes

Short- and medium-chain alkanes, which can be used as transportation fuel, may be produced by genetically modified lipid-producing microalgae. This conversion involves transforming fatty acids into aldehydes and then into alkanes, with the final step potentially catalyzed by a decarbonylase enzyme, though no functional decarbonylase has been cloned yet (Gui et al., 2021). The green microalga *B. braunii*, known for producing very-long-chain alkanes, has been studied for this process. Different strains of *B. braunii* produce varying long-chain hydrocarbons, with strain A producing dienes and trienes, and strain B producing triterpenoid hydrocarbons (Phummala et al., 2014). Decarbonylase activity has been found in the leaves of *Pisum sativum* and in *A. thaliana*, where enzymes like Cer1 and Cer22 are thought to be involved in wax formation. The alkanes produced by these decarbonylases are typically very long-chain (22+ carbons) and require further processing for fuel (Hall and Payne, 1997). Some long-chain alkane production has been reported in *Vibrio furnissii*, though later studies have questioned these results. Producing shorter-chain alkanes suitable for direct fuel use remains an ongoing goal, requiring further research to understand the enzymes involved (Kraan, 2012).

6. Genetic Engineering of Microalgae to Increase Lipid Production

Microalgae have low energy density due to their high water content and relatively low levels of proteins and lipids, resulting in higher production costs. Genetic engineering can reduce these costs by 15–20% (Chong et al., 2021). Most studies focus on modifying genes related to lipid metabolism, either by knocking out or introducing genes. For example, overexpression of the enzyme NeodGAT2

in *N. oleoabundans* increases TAG content, while knocking out the phospholipase A2 gene in *C. reinhardtii* boosts lipid production by 64.25% (Huang et al., 2016). Cyanobacteria face stress in biofuel synthesis, so genetic engineering efforts focus on enhancing biofuel compound transfer, like overexpressing thioesterase in *Synechocystis sp.* to increase FA secretion (Hewelt-Belka et al., 2016). Genetic engineering is advancing microalgae biofuel production, but requires a deep understanding of metabolic pathways (Fawzy et al., 2021). Omics and genetic tools are keys for reducing production costs and improving efficiency.

6.1 Types of Genetic Engineering Resources in Microalgae

6.1.1 Algomics resources

Omics techniques, collectively known as ‘algomics’ (including genomics, transcriptomics, proteomics, metabolomics, and metagenomics), play a crucial role in microalgae research, helping design experiments and predict outcomes related to lipid production (Hui et al., 2016). The integration of these methods with bioinformatics aids in developing economically viable microalgal biofuels and discovering new species. Genome sequencing has advanced, with projects like 10KP aiming to sequence 1000 green algae genomes (Fuad-Hossain et al., 2020). Genome editing, including gene knockout, is used to enhance biofuel production. Transcriptomics reveals gene expression related to lipid production, while proteomics focuses on protein regulation, identifying key proteins for lipid enhancement (Gui et al., 2021). Metabolomics analyzes low molecular compounds and helps optimize lipid biosynthesis. Combined multi-omics approaches provide a comprehensive understanding of microalgal metabolism, facilitating the control of production pathways and improving biofuel yields (Kraan, 2012).

6.1.2 Mutant Resources

Forward genetics, using random mutagenesis and adaptive laboratory evolution, can generate mutant phenotypes without prior knowledge of an organism’s genetics. Methods like γ -ray, UV exposure, and chemical mutagenesis have been used to create high lipid-accumulating mutants in species like *N. gaditana*, *N. oculata*, and *Scenedesmus sp.* (Hewelt-Belka et al., 2020). However, random mutagenesis can be non-specific, unstable, and harmful, requiring careful selection methods. Adaptive laboratory evolution subjects microalgae to selective pressures over time to improve traits (Krishnan et al., 2015). Insertional mutagenesis integrates foreign DNA into the microalgal genome to direct metabolism toward lipid synthesis. While *Chlamydomonas reinhardtii* face challenges in transformation efficiency and identification of insertion site being useful for studying gene function and mutant libraries at the same time (Kumar et al., 2020).

6.2 Genetic Modification Techniques in Microalgae for Biofuel Production

6.2.1 Procedures of Transformation

The technique of genetic engineering includes introducing foreign DNA into the genome, and ensuring stable expression thus transforming microalgal cells. It requires careful selection of methods depends on the species and the type of modification to increase transformation efficiency (Nagappan et al., 2020). There are multiple techniques such as electroporation, particle bombardment, sonication, viruses, and agrobacterium-mediated transformation. To increase lipid content in microalgae, techniques like electroporation and particle bombardment offer the highest efficiency, even without removing cell walls (Kumar et al., 2016). However, for lipid accumulation a higher rate of false positives and less effective results are shown by agrobacterium-mediated transformation (Popovich et al., 2019). There are also some less widely used methods like sonication and silicon carbide whiskers.

6.2.2 Power of Genome Editing Tools

To improve product quality and quantity in microalgae novel genome-editing tools like transcription activator-like effector nucleases (TALENs), zinc-finger nucleases (ZFNs), RNA interference (RNAi) and CRISPR/Cas9 have been used (Roy-Chong et al., 2022). In *P. tricornutum* and *T. pseudonana*, lipid content is enhanced through silencing genes by using RNAi (Lv et al., 2019). For lipid accumulation in *C. reinhardtii*, ZFNs have also been used. Bounded by hard design and off-target effects lipid production can be increased by TALENs. So for more efficient and accurate results, CRISPR/Cas9 uses guide RNA to create precise DNA breaks which is applied to increase lipid accumulation in microalgal strains such as *C. reinhardtii*, *N. oceanica*, *N. gaditana*, and *P. tricornutum* (Santana-Sanchez et al., 2021).

6.2.3 Expression of Foreign Gene in Microalgal Strains

To manufacture biofuels, such as biogas, ethanol, or biodiesel, there is a technique that employs inserting genes from other microbes into algal strains to improve their ability, called foreign gene expression. For the making of biofuels, enhancing the growth rate, lipid production, and other cellular processes occurs through this method (Mairet et al., 2011). Transgenic procedures like: Agrobacterium-mediated gene transfer method uses *Agrobacterium tumefaciens* to transfer foreign DNA into algal cells. The inserted genes can help to enhance lipid production or improve the algae's ability to convert sunlight into expandable energy. Another method involves applying an electric field to algae cells, which temporarily modify the cell membrane, allowing foreign DNA to enter which is called electroporation (Maliha and Abu-Hijleh, 2022). This is useful for algal strains that are complex to transform using other methods. Another method called gene gun uses high-velocity gold or tungsten particles coated with DNA to shoot

the DNA into the algae cells. This is widely used for species like *Chlamydomonas reinhardtii*. Genetic manipulations can be made to enhance cellular pathways involved in biofuel production. For example, by increasing levels of triacylglycerols (TAGs) through inserting genes that enables algae to make biodiesel (Bra'nyikova et al., 2011). To maximize biofuel output there are some common genes inserted to include those encoding enzymes from other organisms that increase the production of sugars, oils, or even genes that redirect carbon metabolism. To enhance the metabolic processes that lead to biofuel accumulation there is a technique that allows for knocking out genes that is CRISPR-Cas9 which is a powerful tool for making precise edits to the algae genome and has been used to improve stress tolerance and lipid production in algae (Metting, 1996). In addition to inserting foreign genes, researchers use specific promoters and enhancer elements to drive the expression of these genes under optimal conditions (Yamada and Sakaguchi, 1982). For example, during periods of high light intensity using light-regulated promoters is an innovative approach to increase biofuel production. So to produce biofuels more efficiently, algae can be engineered by integrating synthetic biology tools. It includes creating synthetic pathways that traditionally not occurs in algae, such as pathways for isoprenoid-based biofuels (like isopropanol or biodiesel). To help the algae to adapt in environmental stresses like: nutrient limitation and temperature fluctuations, synthetic biology permits for the making of innovative metabolic circuits while maximizing biofuel output (Nagappan et al., 2020). To convert biomass into biofuels more efficiently, algae is co-culturing with engineered bacteria or fungi which involves genetic modifications in bacteria to increase the breakdown of algae-derived materials. For biofuel production algae's resistance to environmental stress can be improved through Genetic modifications for increasing overall productivity (Metting, 1996). For example, to improve salt tolerance or boost photosynthetic efficiency, certain genes can be incorporated in algae to produce more biomass or lipids for biofuel production.

6.2.4 Silencing of Gene Through Rna Interference

To increase biofuel production in microalgae there is a technique for silencing or down regulating specific genes called RNA interference (RNAi) which is a powerful tool, and it has been effectively applied to regulate genes involved in lipid synthesis, stress response, or metabolic pathways that affect algal growth and biofuel accumulation (Nakanishi et al., 2014). The RNAi mechanism works by incorporating small RNA molecules such as small interfering RNA or siRNA that target and destroy messenger RNA (mRNA), preventing the expression of specific genes (Hall and Payne, 1997).

6.2.4.1 Significant Uses of RNA Interference in Microalgae for Biofuel Production

6.2.4.1.1 Boosting up Lipid Accumulation for production of Biodiesel

It has been seen that for efficient biodiesel, microalgae can produce lipids like triglycerides by silencing genes included in lipid breakdown or promoting lipid

biosynthesis pathways which can be increased in microalgae cells by using RNAi. (Noraini et al., 2014). To make algae more suitable for biodiesel production, silencing of genes like lipase which catabolize lipids and acetyl-CoA carboxylase which is involved in fatty acid biosynthesis can be done (Mairet et al., 2011).

6.2.4.1.2 Regulation of Photosynthesis and Fixation of Carbon

RNAi can be used to target genes included in carbon fixation which is a critical process for energy production and growth in algae (Phummala et al., 2014). Researchers can potentially redirect carbon flow towards biofuel production by down regulating genes like RuBisCO (ribulose-1,5-bisphosphate carboxylase/oxygenase), which plays a crucial role in the Calvin cycle (Bilad et al., 2012). Apart from that, under certain conditions algal productivity can be increased by silencing genes that limit photosynthesis or inhibit light capture (Ortigueira et al., 2015).

6.2.4.1.3 Modulation in Stress Response

Under stressful conditions like nutrient limitation of nitrogen or phosphorus, high light intensity, or temperature variations algae often produce more amount of lipids or other metabolites (Plohn et al., 2021). RNAi can be used to silence stress-responsive genes that could otherwise inhibit biofuel production. For boosting biomass production and increasing biofuel yields, silencing oxidative stress related genes could enhance the algae's ability to tolerate high-light or high-temperature stresses (Connor and Liao, 2009).

6.2.4.1.4 Stimulating the Pathways for Synthesis of Biofuels

It has been observed that to favor lipid accumulation, pathways that lead to the production of starch or other carbohydrates can be down regulated through RNAi technology (Popovich et al., 2019). Additionally, the overall yield of biofuels can be enhanced by shifting metabolic fluxes towards biofuel-compatible molecules through silencing genes involved in fatty acid modification or degradation (Nakanishi et al., 2014).

6.2.4.1.5 Profile Improvement of Fatty Acid

For biodiesel production, the alteration in fatty acid profile of microalgae to produce more desirable fatty acids can be done by RNAi (Roy et al., 2014). The algal strains can be engineered to produce higher proportion of unsaturated fatty acids to improve the fuel's quality which can be achieved by silencing or over expressing certain enzymes in the fatty acid biosynthesis pathway (Roy et al., 2022). Apart from that shutdown genes that negatively affect growth and biomass accumulation like genes involved in regulation of cell cycle or stress responses can lead to high level of algal biomass production which directly increases potential feedstock for biofuel production. (Dubini et al., 2009).

6.2.4.1.6 Successful Examples in Algal Strains

- ***Chlamydomonas reinhardtii***: *C. reinhardtii* is a broadly studied model organism in biofuel research, It has been modified using RNAi to enhance production of lipid. For instance, the decrease in gene expression involved in biosynthesis of starch like: ADP-glucose pyrophosphorylase can affect carbon flux towards accumulation of lipids (Krishnan et al., 2015).
- ***Nannochloropsis species***: Due to the high lipid content in *Nannochloropsis* it is another microalga commonly used for biodiesel production. To increase lipid content certain genes involved in lipid degradation or starch synthesis has been silenced by using RNAi for biofuel production (Yanlin et al., 2014).
- ***Phaeodactylum tricornutum***: By regulating lipid metabolism and stress tolerance this algal species has also been genetically modified with RNAi to improve biofuel yields. Scientists have switched off genes like acetyl-CoA carboxylase, which controls biosynthesis of fatty acid (Bagchi and Mallick, 2016).

6.3 The antisense RNA Technology

This technique involves incorporating a complementary RNA strand to the mRNA of gene of interest. To prevent translation and effective silencing of the genes, the antisense RNA binds to the mRNA (Santana-Sanchez et al., 2021). To favor lipid biosynthesis metabolic pathways are shifted by down regulating genes involved in starch production or carbon storage by using antisense RNA (Saini et al., 2021). For higher lipid accumulation the inhibition of genes involved in degradation of fatty acid can be done by this technique. There is a species of microalga called *Nannochloropsis* which has high lipid content, but antisense RNA strategies has been targeted by silencing genes involved in starch synthesis to increase lipid production (Shin et al., 2018).

6.4 Transcriptional Gene Silencing (TGS)

This technique involves long-term gene silencing by manipulating the structure of chromatin which is done through DNA methylation or histone modification and prevents transcription (Shiels et al., 2021). To get high-yield biofuel producing microalgal strains TGS can be used to silence genes permanently that inversely affect lipid accumulation or biomass production (Saravanan et al., 2022). Also under certain environmental stresses, like high light intensity or nutrient starvation TGS can also be applied to turnoff genes that bounds algal growth or synthesis of biofuel. This technique has been experimented as a candidate method for enhancing stress tolerance and improving production of biofuel by silencing regulatory genes (Artamonova et al., 2017).

6.5 Virus-mediated Gene Silencing

The bacterial virus (phage)-based systems such as some viral vectors have been chosen for gene silencing in algae. This can be done through RNAi

mechanisms where viral vectors deliver RNA or DNA fragments into algal cells to induce gene silencing (Shin et al., 2018). On the industrial scale viral-mediated gene silencing emerges as a useful tool for large-scale gene silencing, helping to improve biofuel production in algal cultures. There are certain genes responsible for lipid degradation which can be silenced through this technique to enhance algal biofuel synthesis (Srinivasan et al., 2018).

6.6 Post-transcriptional Gene Silencing (PTGS)

It is a process that occurs prior the translation into protein and after the mRNA is transcribed so it is called Post-transcriptional gene silencing (PTGS) (Srinivasan et al., 2018). It involves the degradation of target mRNA to inhibit translation by using small RNA molecules, similar to RNAi. To optimize the metabolism of algae for biofuel production PTGS can be used to silence genes involved in lipid degradation or carbon flux regulation. Faster growth rates and improved lipid accumulation in microalgae can also be achieved by silencing certain regulatory genes using PTGS (Huang et al., 2016).

6.7 Gene disruption

6.7.1 TALENs (Transcription Activator-Like Effector Nucleases)

This technique induces double-strand breaks at specific genomic locations by using engineered nucleases. TALENs work identical to CRISPR-Cas9, but they depend on a different procedure for identifying DNA sequences (Subramani and Venu, 2019). To redirecting resources towards lipid production, disruption of genes that lead to the production of non-oil metabolites and other non-essential pathways can be done by using TALENs. The improvement in algal growth and lipid production can be achieved by disrupting genes involved in starch synthesis and induce algae to focus more energy toward lipid accumulation (Ullah et al., 2014). It has been observed under nitrogen-limited conditions that TALENs have been used to knock out pyruvate dehydrogenase kinase in algae, to increase their growth and lipid content, which is positive to production of biofuels (Wieczorek et al., 2014).

6.7.2 Zinc Finger Nucleases (ZFNs)

To incorporate double-strand breaks at specific target sites there is engineered DNA-binding proteins called Zinc finger nucleases that can be used to corrupt genes through homology-directed repair or non-homologous end joining (Uma et al., 2020). To increase suitability for biofuels like biodiesel, ZFNs can be used to interfere genes involved in synthesis or breakdown of carbohydrate, leading to a higher content of lipids produced (Wang et al., 2021). Under certain stress conditions genes that inhibit algal growth are disrupted which can also increase biomass accumulation, leading to better yield of biofuel. In *Chlamydomonas reinhardtii* genes involved in carbon flux have been knocked out by integrating ZFNs to promote lipid accumulation in response to harsh environmental conditions (Ortigueira et al., 2015).

6.7.3 Insertional Mutagenesis

This technique relies on incorporating a selectable marker or a reporter gene at random sites into genome of algae leading to the inactivation of the targeted gene as insertion nullify the function of genes at the site of incorporation (Wang et al., 2010). The metabolic pathways specific to biofuel production are identified by tracing the genes involved in them. A library of mutants can be created using insertional mutagenesis to uplift, stress resistance, improved growth or lipid production under nutrient-deficient conditions (Wieczorek et al., 2014). To favor lipid accumulation genes responsible for carbohydrate or protein biosynthesis can be disrupted. Under nutrient-deficient conditions, random insertional mutagenesis has been used to identify genes that can be corrupted to enhance lipid content or promotion of biomass yield in *Chlamydomonas* and other algal species (Roy et al., 2014).

6.7.4 Shutting Down the Gene Using Homologous Recombination

Homologous recombination permits scientists to change or corrupt a gene by incorporating a fragment that targets the gene of interest for deletion or replacement. This method can be used to make knockouts where the gene is completely shutdown (Yamada and Sakaguchi, 1982). Shutting Non-essential Pathways: By silencing genes included in pathways like starch synthesis, more metabolic energy is focused towards biofuel-precursor production (Yanlin et al., 2014). Under limited growth conditions homologous recombination successfully disabled genes for pyruvate dehydrogenase and acetyl-CoA carboxylase in species like *Chlamydomonas reinhardtii*, to increase lipid synthesis and biomass accumulation (Wang et al., 2010).

6.7.5 Transposon-Mediated Mutagenesis

Transposons are DNA sequences whose insertion can disrupt the function of target genes as they can move from one location in the genome to another (Zhao et al., 2022). This technique can be used to create random malfunctions across the genome, leading to the recognition of useful mutations for biofuel production (Kumar et al., 2016).

6.8 Benefits of Genetic Engineering

Genetic engineering provides a wide variety of uses for improving biomass productivity in algae, creating them more efficient and effective as a candidate of biofuels and other useful compounds (Aizouq et al., 2020). By directly modifying algal genetic material, researchers can improve their growth rates, stress tolerance, nutrient uptake and efficiency. (Cheng et al., 2014).

6.8.1 Enhancement in Growth Rates of Microalgal Biomass

Gene manipulations can magnify algal metabolism, permitting them to grow faster by improving pathways working in energy production, protein synthesis, and carbon fixation, also improving their rates of biomass accumulation (Arora and Philippidis, 2021). Genome engineering can regulate photosynthetic efficiency

by manipulating genes that regulate light-gathering complexes or enzymes of carbon fixation, allowing algae to grasp sunlight and transform it into biomass much efficiently (Bagchi and Mallick, 2016). Also by improving the efficiency of RuBisCO enzyme which is responsible for fixing CO₂ can allow algae to use carbon more effectively, leading to higher biomass production (Phummala et al., 2014).

6.8.2 Improvement in Various Stress Tolerances

Genetic engineering can improve the algal tolerance to fluctuating environmental conditions like temperature changes, nutrient deprivation, high light, or salinity, allowing them to maintain high biomass productivity under non-optimal conditions (Behl et al., 2020). During harsh conditions like oxidative stress or nutrient limitations yield reductions can be prevented by introducing genes that help algae to manage excess light energy like encoding genes for photo protective proteins algae can continue to grow rapidly even under intense sunlight, which is crucial for large-scale outdoor cultivation (Kraan, 2012 and Bergthorsen et al., 2015).

6.8.3 Efficient Nutrient Utilization

Nitrogen and phosphorus are crucial nutrients for algal growth, but they have shortage in the environment. Genetic engineering can be used to improve the efficiency of nutrient uptake, making algae more productive even in nutrient deficient situations (Bharti et al., 2021). For example, manipulating the algal genome to increase the activity of transporters for nitrogen or phosphorus can increase growth rates (Bilad et al., 2012). Algae are often cultivated in regulated environments or photo bioreactors where the availability of nutrients is maintained. Gene manipulations that increase nutrient uptake can nullify the requirement for conventional fertilizers, decreasing costs and enhancing sustainability (Yamada and Sakaguchi, 1982).

6.8.4 Increasing the Lipid content and Biomass Yields

For biofuels like biodiesel, the lipid concentration of algae is important. Genetic engineering can be used to enhance lipid accumulation by modifying genes working in lipid biosynthesis (e.g., over expressing enzymes like acetyl-CoA carboxylase) (Cheng et al., 2014). Greater lipid content transforms directly into higher energy yields per unit of biomass. For specific applications like biofuels, animal feed, or pharmaceuticals gene editing allows for fine-tuning the composition biomass by cahnging carbohydrate, protein, or lipid contents in the algal cells (Bra'nyikova et al., 2011 and Chong et al., 2021).

6.8.5 Increasing the Efficiency of Photosynthesis

Gene manipulations can increase the potential of algae to acheive light energy more efficiently. By regulating the expression of light-harvesting complexes (LHCs), algae can better grasp sunlight, improving the efficiency of photosynthesis and biomass production (Coelho et al., 2014). Genome engineering can enhance the

working of proteins included in photosystem I and II, key parts of photosynthesis, permitting algae to express more efficient light conversion under a wide range of environmental stresses (Conner and Liao, 2009).

6.8.6 Improvement in Fixation of Carbon Dioxide (CO₂)

Genetic engineering can optimize the process of carbon dioxide (CO₂) fixation for algae growth (Dubini et al., 2009). Such as, by improving the function of the carbonic anhydrase enzyme or increasing the functionality of the Calvin cycle, algae can grow faster and gather more biomass in CO₂-rich conditions (Fawzy et al., 2021). By genetically improving algae's potential to tolerate high concentrations of CO₂, they can be cultured in conditions where CO₂ is more accessible, such as in carbon capture systems or industrial flue gas emissions. This could lead to provide a solution for CO₂ mitigation and higher biomass productivity (Metting, 1996).

6.8.7 Rapid Cultivation Cycles

Genome editing can alter the reproduction cycle of algae, making to higher cultivation cycles (Fuad-Hossain et al., 2020). This is significantly useful for algae used in biofuel production, where rapid turnover and high productivity are crucial to make biofuels economically feasible. Genome manipulations can balance growth phases, allowing algae to be collected at peak productivity, thereby increasing biomass yields over lesser time durations (Hall and Payne, 1997).

6.8.8 Greater stability of Genetic Makeup

Genetically engineered algae can be designed to have stable, heritable traits that increase biomass productivity. This stability allows for the development of high-yield algal strains that can be cultivated consistently over time without losing desirable traits, providing a reliable source of biomass (Gui et al., 2021). Some genetic engineering strategies, such as using marker genes or selective pressure techniques, can reduce genetic drift in algae populations, maintaining high biomass productivity over successive generations (Mairet et al., 2011).

6.8.9 Boosting the Performance of Bioreactors

Genetic engineering can also optimize algae for growth in bioreactors, such as photobioreactors or fermenters (Hall and Payne, 1997). Traits like increased resistance to contamination, improved nutrient utilization, and better light utilization can result in more efficient bioreactor operation, leading to higher productivity (Hewelt-Belka et al., 2016). By modifying algae to grow more efficiently in different media (e.g., wastewater, CO₂-enriched air, or saline water), genetic engineering can open up new possibilities for large-scale biomass production that uses waste resources or non-potable water (Bilad et al., 2012).

6.8.10 Contamination Hazards Reduction

Genetic engineering can introduce mechanisms that make algae more resistant to common contaminants, such as bacteria, fungi, or other microorganisms (Huang

et al., 2016). This can ensure that algae cultures remain healthy and productive, reducing the need for external inputs like antibiotics and chemical treatments (Kraan et al., 2012). Genetically engineered algae that are more resistant to contamination are better suited to cultivation in closed systems, making industrial-scale algae biofuel production more reliable and cost-effective (Connor and Liao, 2009).

6.9 Shortcomings and Hazards in Genetic Engineering of Microalgae

Genetic engineering in microalgae has significant potential for boosting lipid production, but concerns arise over the environmental and ethical impacts of GM microalgae releases (Krishnan et al., 2015). These organisms could escape into the environment via wind, water, or accidents, causing unintended ecological consequences such as disruption of food webs, displacement of native species, or harmful algal blooms (Saravanan et al., 2022). To mitigate risks, regulations and insecurity measures are essential, and developing “suicide genes” to control hazardous strains is recommended. Additionally, challenges exist in scaling transgenic microalgae for large-scale cultivation, as laboratory conditions may not replicate well in open environments, impacting growth rates, gene quality, and productivity. No outdoor cultivation of GM microalgae has been reported due to these risks (Shiels et al., 2021).

7. Insight to the Future

Significant progress has been made in selecting and optimizing lipid-producing microalgal strains, but further improvements are needed for microalgal biofuels to compete with fossil fuels. Advances in dehydration, harvesting techniques, genome sequencing, gene editing, and synthetic biology will support the development of microalgae as a renewable energy source (Subramani and Venu 2019). Progress in lipid extraction, purification techniques, and AI applications in extraction processes are also essential. Additionally, exploring the simultaneous production of biofuels and high-value byproducts holds promise (Uma et al., 2020 and Zhao et al., 2022). The future of microalgal biofuels relies on cost-effective methods and efficient technologies to accelerate commercialization. This discussion primarily focuses on enhancing lipid accumulation in eukaryotic microalgae, leaving out details on harvesting, extraction, and purification processes (Kumar et al., 2020).

8. Summary

To address the crisis, microalgal biofuels offer a promising solution to fossil fuels, but challenges remain a barrier to achieving cost-efficient large-scale production. Selecting resilient and adaptable microalgae strains is the first step. To influence yield and performance, gene editing and multi-omics technologies provide more precise production methods, controlling cultivation conditions and adjusting photobioreactor (PBR) parameters. However, careful consideration of environmental and human health risks associated with large-scale outdoor cultivation of genetically modified microalgae is required. Other developed countries like china faces observable challenges in boosting up biofuel production

from microalgae, but a promising future is underway with the synergy between advancing technologies and market demands. In few years, the risks associated with genetically modified will be overcome by exploring synthetic biology which could play a key role in mitigating and improving the economic feasibility of biofuel production.

References

- Aizouq, M., Peisker, H., Gutbrod, K., Melzer, M., Holzl, G. and Dormann, P. (2020). Triacylglycerol and phytol ester synthesis in *Synechocystis* sp. PCC6803. *Proceed Nat Acad Sci*, 117(11):6216–22.
- Arora, N. and Philippidis, G.P. (2021). Insights into the physiology of *Chlorella vulgaris* cultivated in sweet sorghum bagasse hydrolysate for sustainable algal biomass and lipid production. *Sci Rep*, 11(1):6779.
- Artamonova, E. Y., Vasskog, T. and Eilertsen, H. C. (2017). Lipid content and fatty acid composition of *Porosira glacialis* and *Attheya longicornis* in response to carbon dioxide (CO₂) aeration. *PLoS ONE*, 12(5): e0177703.
- Bagchi, S. and Mallick, N. (2016). Carbon dioxide biofixation and lipid accumulation potential of an indigenous microalga *Scenedesmus obliquus* (Turpin) Kutzing GA 45 for biodiesel production. *RSC Adv*, 6:29889–98.
- Behl, K., Sessa-Charan, P., Joshi, M., Sharma, M., Mathur, A., Kareya, M.S., Jutur, P.P., Bhatnagar, A. and Nigam, S. (2020). Multifaceted applications of isolated microalgae *Chlamydomonas* sp. TRC-1 in wastewater remediation, lipid production and bioelectricity generation. *Bioresour Technol*, 304:122993.
- Bergthorsen, J. M. and Thomson, M. J. (2015). A review of the combustion and emissions properties of advanced transportation biofuels and their impact on existing and future engines, *Renewable Sustainable Energy Rev*, 42: 1393–1417.
- Bharti, M.K., Chalia, S., Thakur, P., Sridhara, S.N., Thakur, A. and Sharma, P.B. (2021). Nanoferrites heterogeneous catalysts for biodiesel production from soybean and canola oil: a review. *Environ Chem Lett*, 19(5):3727–46.
- Bilad, M. R., Vandamme, D., Foubert, I., Muylaert, K. and Vankelecom, I. F. (2012). Harvesting microalgal biomass using submerged microfiltration membranes. *Bioresour Technol*, 111:343–52 BP Statistical Review of World Energy. 2024.
- Bra'nyikova', I., Mars 'al'kova', B., et al., (2011). Microalgae—novel highly efficient starch producers. *Biotechnol. Bioeng*, 108 (4), 766–776.
- Cheng, J., Liu, Y. Q., Lin, R. C., Xia, A., Zhou, J. H. and Cen, K. F. (2014). Cogeneration of hydrogen and methane from the pretreated biomass of algae bloom in Taihu Lake. *Int J Hydrogen Energy*, 39(33): 18793–802.
- Chong, J.W.R., Khoo, K.S., Chew, K.W., Ting, H.Y. and Show, P.L. (2023). Trends in digital image processing of isolated microalgae by incorporating classification algorithm. *Biotechnol Adv*, 63: 108095.
- Chong, J.W.R., Khoo, K.S., Yew, G.Y., Leong, W.H., Lim, J.W., Lam, M.K., Ho, Y.C., Ng, H.S., Munawaroh, H.S.H. and Show, P.L. (2021). Advances in production of bioplastics by microalgae using food waste hydrolysate and wastewater: a review. *Bioresour Technol*, 342: 125947.
- Chong, J.W.R., Yew, G.Y., Khoo, K.S., Ho, S.H. and Show, P.L. (2021). Recent advances on food waste pretreatment technology via microalgae for source of polyhydroxyalkanoates. *J Environ Manage*, 293: 112782.
- Coelho, M. S., Barbosa, F. G., Zimmerman de Souza, M. D. R. A. (2014). The scientometric research of microalgal biomass as a source of biofuel feedstock. *Algal Res*, 6 (Part B):132–138.

- Connor, M. R. and Liao, J. C. (2009). Microbial production of advanced transportation fuels in non-natural hosts. *Curr. Opin. Biotechnol*, 20: 307–315.
- Dubini, A., Mus, F., Seibert, M., Grossman, A. R. and Posewitz, M. C. (2009). Flexibility in anaerobic metabolism as revealed in a mutant of *Chlamydomonas reinhardtii* lacking hydrogenase activity. *J. Biol. Chem.* 284:7201–7213
- Fawzy, M.A., El-Otify, A.M., Adam, M.S. and Moustafa, S.S.A. (2021). The impact of abiotic factors on the growth and lipid accumulation of some green microalgae for sustainable biodiesel production. *Environ Sci Pollut Res*, 28(31):42547–61.
- Fuad-Hossain, M., Ratnayake, R.R., Mahbub, S., Kumara, K.L.W. and Magana-Arachchi, D.N. (2020). Identification and culturing of cyanobacteria isolated from freshwater bodies of Sri Lanka for biodiesel production. *Saudi J Biol Sci*, 27(6):1514–20.
- Garcia-Moscosa, J. L., Obeid, W., Sandeep, K., et al. (2013). Flash hydrolysis of microalgae (*Scenedesmus* sp.) for protein extraction and production of biofuels intermediates. *J Supercrit Fluids*, 82:183–190.
- Gui, J., Chen, S., Luo, G., Wu, Z., Fan, Y., Yao, L. and Xu, H. (2021). Nutrient deficiency and an algicidal bacterium improved the lipid profiles of a novel promising oleaginous dinoflagellate, *Prorocentrumdonghaiense*, for biodiesel production. *Appl Environ Microbiol*, 87(19):e01159-e11121.
- Hall, J. and Payne, G. (1997). Factors controlling the growth of field population of *Hydrodictyonreticulatum* in New Zealand. *Journal of Applied Phycology*, 9: 229-236.
- Hewelt-Belka, W., Kot-Wasik, A., Tamagnini, P. and Oliveira, P. (2020). Untargeted lipidomics analysis of the cyanobacterium *Synechocystis* sp. PCC 6803: lipid composition variation in response to alternative cultivation setups and to gene deletion. *Int J Mol Sci*, 21(23):8883.
- Huang, J. J., Bunjamin, G., Teo, E. S., Ng, D. B. and Lee, Y. K. (2016). An enclosed rotating floating photobioreactor (RFP) powered by flowing water for mass cultivation of photosynthetic microalgae. *Biotechnol Biofuels*, 9(1):218.
- Hui, W., Wenjun, Z., Wentao, C., Lili, G. and Tianzhong, L. (2016). Strategy study on enhancing lipid productivity of filamentous oleaginous microalgae *Tribonema*. *Bioresour Technol*, 218:161–6.
- Kraan, S. (2012). Algal polysaccharides, novel applications and Outlook, *Carbohydrates–comprehensive studies on glycobiology and glycotecnology*, 978-953.
- Krishnan, V., Uemura, Y., Thanh, N. T., Khalid, N. A., Osman, N. and Mansor, N. (2015). Three types of Marine microalgae and *Nannochloropsisoculata*cultivation for potential source of biomass production. *J Phys: Conf Ser*, 622: 012034.
- Kumar, G., Sen, B., Sivagurunathan, P. and Lin, C. (2016). High rate hydrogen fermentation of cello-lignin fraction in de-oiled jatropha waste using hybrid immobilized cell system. *Fuel*, 182: 131–40.
- Kumar, G., Shekh, A., Jakhu, S., Sharma, Y. and Sharma, T.R. (2020). Bioengineering of microalgae: recent advances, perspectives, and regulatory challenges for industrial application. *Front Bioeng Biotechnol*, 8:914.
- Lv, Z., Chu, A.M.Y., McAleer, M. and Wong, W.K. (2019). Modelling economic growth, carbon emissions and fossil fuel consumption in china: co-integration and multivariate causality. *Int J Environ Res Public Health*, 16(21):4176.
- Mairet, F., Bernard, O., Masci, P, Lacour, T. and Sciandra, A. (2011). Modelling neutral lipid production by the microalga *Isochrysisaff. galbana* under nitrogen limitation. *Bioresour Technol*, 102:142–9.
- Maliha, A. and Abu-Hijleh B. (2022). A review on the current status and post pandemic prospects of third-generation biofuels. *Energy Syst*, 14:1–32.
- Metting, F. B. (1996). Biodiversity and application of microalgae, *J. Ind. Microbiol. Biotechnol*, 17 477–489.

- Nagappan, S., Bhosale, R., Duc-Nguyen, D., Pugazhendhi, A., Tsai, P.C., Chang, S.W., Ponnusamy, V.K. and Kumar, G. (2020). Nitrogen-fixing cyanobacteria as a potential resource for efficient biodiesel production. *Fuel*, 279: 118440.
- Nakanishi, A., Aikawa, S., Ho, S. H., Chen, C. Y., Chang, J. S., Hasunuma, T. and Kondo, A. (2014). Development of lipid productivities under different CO₂ conditions of marine microalgae *Chlamydomonas* sp. JSC4. *Bioresour Technol*, 152:247–52.
- Noraini, M. Y., Ong, H. C., Badrui, M. J, et al. (2014). A review on potential enzymatic reaction for biofuel production from algae. *Renewable Sustainable Energy Rev*, 39: 24–34.
- Ortigueira, J., Alves, L., Gouveia, L. and Moura, P. (2015). Third generation biohydrogen production by *Clostridium butyricum* and adapted mixed cultures from *Scenedesmus obliquus* microalga biomass. *Fuel*, 153: 128–34.
- Phummala, K., Imai, T., Reungsang, A., Chairattananokorn, P., Sekine, M., Higuchi, T., Yamamoto, K. and Kanno, A. (2014). Delignification of disposable wooden chopsticks waste for fermentative hydrogen production by an enriched culture from a hot spring. *J Environ Sci*, 26: 1361–8.
- Plohn, M., Spain, O., Sirin, S., Silva, M., Escudero-Onate, C., Ferrando-Climent, L., Allahverdiyeva, Y. and Funk, C. (2021). Wastewater treatment by microalgae. *Physiol Plant*, 173:568–78.
- Popovich, C.A., Pistonesi, M., Hegel, P., Constenla, D., Bielsa, G.B., Martin, L.A., Damiani, M.C. and Leonardi, P.I. (2019). Unconventional alternative biofuels: quality assessment of biodiesel and its blends from marine diatom *Naviculacincta*. *Algal Res*, 39: 101438.
- Roy, S., Kumar, K., Ghosh, S. and Das, D. (2014). Thermophilic biohydrogen production using pre-treated algal biomass as substrate. *Biomass Bioenergy*, 61: 157–66.
- Roy-Chong, J.W., Tan, X., Khoo, K.S., Ng, H.S., Jonglertjunya, W., Yew, G.Y. and Show, P.L. (2022). Microalgae-based bioplastics: future solution towards mitigation of plastic wastes. *Environ Res*, 206: 112620.
- Saini, R., Osorio-Gonzalez, C.S., Brar, S.K. and Kwong, R. (2021). A critical insight into the development, regulation and future prospects of biofuels in Canada. *Bioengineered*, 12(2):9847–59.
- Santana-Sanchez, A., Lynch, F., Sirin, S. and Allahverdiyeva, Y. (2021). Nordic cyanobacterial and algal lipids: Triacylglycerol accumulation, chemotaxonomy and bioindustrial potential. *Physiol Plant*, 173(2):591–602.
- Saravanan, A., Deivayanai, V.C., Senthil-Kumar, P., Rangasamy, G. and Varjani, S. (2022). CO₂ bio-mitigation using genetically modified algae and biofuel production towards a carbon net-zero society. *Bioresour Technol*, 363: 127982.
- Shiels, K., Tsoupras, A., Lordan, R., Nasopoulou, C., Zabetakis, I., Murray, P. and Saha, S.K. (2021). Bioactive lipids of marine microalga *Chlorococcum* sp. SABC 012504 with anti-inflammatory and anti-thrombotic activities. *Mar Drugs*, 19(1):28.
- Shin, Y. S., Choi, H. I., Choi, J. W., Lee, J. S., Sung, Y. J. and Sim SJ. (2018). Multilateral approach on enhancing economic viability of lipid production from microalgae: a review. *Bioresour Technol*, 258:335–44.
- Srinivasan, R., Mageswari, A., Subramanian, P., Suganthi, C., Chaitanyakumar, A., Aswini, V. Gothandam, K. M. (2018). Bicarbonate supplementation enhances growth and biochemical composition of *Dunaliella salina* V-101 by reducing oxidative stress induced during macronutrient deficit conditions. *Sci Rep*, 8(1):6972.
- Subramani, L. and Venu, H. (2019). Evaluation of methyl ester derived from novel *Chlorella emersonii* as an alternative feedstock for DI diesel engine & its combustion, performance and tailpipe emissions. *Heat Mass Transfer*, 55(5):1513–34.

- Ullah, K., Ahmad, M., Sofia, et al. (2014). Algal biomass as a global source of transport fuels overview and development perspectives. *Progress Nat Sci Mat Int*, 24 (4):329– 339.
- Uma, V.S., Gnanasekaran, D., Lakshmanan, U. and Dharmar, P. (2020). Survey and isolation of marine cyanobacteria from eastern coast of India as a biodiesel feedstock. *Biocatal Agric Biotechnol*, 24: 101541.
- Wang, M., Liu, H., Qiao, K., Ye, X., Takano, T., Liu, S. and Bu, Y. (2021). Exogenous NaHCO₃ enhances growth and lipid accumulation of the highly NaHCO₃-tolerant *Nannochloris* sp. JB17. *J Appl Phycol*, 33:241–53.
- Wang, Y., Wang, H., Feng, X. Q., Wang, X. F. and Huang, J. X. (2010). Biohydrogen production from cornstalk wastes by anaerobic fermentation with activated sludge. *Int J Hydrogen Energy*, 35 (7):3092–9.
- Wieczorek, N., Kucuker, M.A. and Kuchta, K. (2014). Fermentative hydrogen and methane production from microalgal biomass (*Chlorella vulgaris*) in a two-stage combined process. *Appl Energy*, 132:108–17
- Yamada, T. and Sakaguchi, K. (1982). Comparative studies on *Chlorella* cell walls – induction of protoplast formation, *Arch. Microbiol*, 132 10–13.
- Yanlin, Z., Cheng, Y., Junhan, L. and Guangrong, H. (2014). Lipid production by a CO₂- tolerant green microalga, *Chlorella* sp. MRA-1. *J Microbiol Biotechnol*, 24(5):683–9.
- Zhao, K., Li, Y., Yan, H., Hu, Q. and Han, D. (2022). Regulation of light spectra on cell division of the unicellular green alga *Haematococcus pluvialis*: insights from physiological and lipidomic analysis. *Cells*, 11(12):1956.

BIOFUELS PRODUCTION USING SUSTAINABLE BIOPROCESSING TECHNOLOGIES

VOL 4

The book sections cover the thermodynamics and kinetics of enzyme-based fuel cells, bridging scientific principles with real-world applications in bioenergy. This book serves as an essential resource for researchers, engineers, and policymakers interested in the advancement of sustainable biofuels. It offers a well-rounded perspective, combining innovations in biofuel production with waste management strategies and bioprocessing techniques to support a greener energy future.

In Chapter 1, we start with an Algal Biofuel Utilization: Challenges and Prospects. Chapter 2 looks at how Production of Bioenergy using Algal Biomasses: Challenges and Prospects. Chapter 3 focuses on how the Exploring the Biofuel Production Through Algal Genetic Engineering: Current Techniques and Challenges, while Chapter 4 highlights the role of Smart Energy Management and Recovery Towards Sustainable Bioenergy, Chapter 5 introduces the Pre-Treatment of Lignocellulosic Wastes for Biofuel Production. Chapter 6 Valorization of agricultural wastes for Glycerol production. Chapter 7 explains how Valorization of Agricultural Wastes for Bioethanol Production, Chapter 8 discusses how Smart fuel and its Management Towards Sustainable Bioenergy, Chapter 9 looks at how Cellulosic Biomass Fermentation for Bioenergy Generation, Chapter 10 explores how Lignocellulosic biomass biorefineries towards circular economy, Chapter 11 explains how Advances in Green Microbial Biofuel Production and Chapter 12 focuses on Lignocellulosic Biomass for the Production Biohydrogen.

This book aims to simplify complex ideas and provide practical insights into how Biomass can revolutionize biofuels, helping to meet the world's growing energy needs in an environmentally friendly way.



**NIPA GENX ELECTRONIC
RESOURCES & SOLUTIONS P. LTD.**

101,103, Vikas Surya Plaza, CU Block
L.S.C.Market, Pitam Pura, New Delhi-110 034
Ph : +91 11 4386 0225, 9717133558, 9540816132
E-mail: newindiapublishingagency@gmail.com
Website: www.niparesources.com

