



आईएफटीएम विश्वविद्यालय, मुरादाबाद, उत्तर प्रदेश

IFTM University, Moradabad, Uttar Pradesh

NAAC ACCREDITED

E-Content

IFTM University, Moradabad

TOPIC: REPRODUCTIVE PHYSIOLOGY

The Male Reproductive System

By:

Mr. Munna Singh


M. Pharm (Pharmacology)

School of Pharmaceutical Sciences IFTM

University, Moradabad.

2023

Sex Determination: Overview




Sexual Differentiation: Internal Embryonic Development

- Bipotential tissues: genes & hormones direct differentiation
- Gonad → testis or ovary
- Wolffian duct → Vas deferens,
- Mullerian duct → oviduct


Sexual Differentiation: Internal Embryonic Development


(a) Development of internal organs



Sexual Differentiation: External Genitalia

(b) Development of external genitalia






Regulation of Reproduction: General Pathways

- **Hypothalamus:** pulse generator
 - Gonadotropin releasing H (GnRH)
- **Anterior Pituitary**
 - Lutenizing H (LH)
 - Follicle stimulating H (FSH)
- **Ovary:**
 - Estrogen, progesterone, Inhibin
- **Testis:** testosterone


Regulation of Reproduction: General Pathways




Male Reproductive Anatomy and Physiology

- Testis
- Epididymis
- Vas deferens
- Seminal vesicle
- Prostate
- Bulbourethral
- Ejaculatory duct
- Urethra
- Penis



Male Reproductive Anatomy




The Male Reproductive System in Anterior View



The Seminiferous Tubules



The Seminiferous Tubules



(c)


Sertoli Cells

- **Form blood-testes barrier:**
 - Prevents autoimmune destruction of sperm.
 - Produce FAS ligand which binds to the FAS receptor on surface to T lymphocytes, triggering apoptosis of T lymphocytes.
 - Prevents immune attack.
- **Secrete inhibin.**
- **Phagocytize residual bodies:**
 - May transmit information molecules from germ cells to Sertoli cells.
- **Secrete androgen-binding protein (ABP):**
 - Binds to testosterone and concentrates testosterone in the tubules.

Spermatogenesis

- Seminiferous tubules
 - Contain spermatogonia
 - Stem cells involved in spermatogenesis
 - Contain Sertoli (sustentacular) cells
 - Sustain and promote development of sperm


Spermatogenesis




Spermiogenesis

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.

- Maturation of spermatozoa.
- Phagocytosis of cytoplasm by the Sertoli cells.
 - Cytoplasm is eliminated.




Spermiogenesis and Spermatozoon Structure



Regulation of Spermatogenesis

- GnRH → LH → Leydig cells → testosterone → growth and division of germ cells
- GnRH → FSH → Sertoli cells → spermatocyte maturation
- Inhibin feedback – FSH, testosterone – short & long loops
- Estrogen
- Growth hormone

Regulation of Spermatogenesis



Maturation of Sperm in Epididymis

- Sperms in the early portion of epididymis are nonmotile
- After 18-24 h they develop *capability of motility*
- Most of sperms are stored in epididymis
- After ejaculation they become motile
- Activity of a sperm is greatly enhanced in neutral to slightly alkaline medium

Seminal Vesicles

- S Vs produce fructose, citric acid and other nutrients as well as prostaglandins and fibrinogen

Prostate

- Slightly alkaline milky fluid that help in neutralizing other seminal fluids as well as the vaginal fluids
- Prostates fluids also contain clotting protein and profibrinolysin

Semen

- Milky white, sticky mixture of sperm and accessory gland secretions (65% of semen is from seminal vesicle, 25% prostate)
- Provides a transport medium and nutrients (fructose), protects and activates sperm, and facilitates their movement
- pH is 7.5
- Prostaglandins in semen:
 - Decrease the viscosity of mucus in the cervix
 - Stimulate reverse peristalsis in the uterus
 - Facilitate the movement of sperm through the female reproductive tract

Semen

- Clotting factors coagulate semen immediately after ejaculation, then fibrinolysin liquefies the sticky mass during the next 15-30 minutes
- After ejaculation, sperms can live 24-48 h
- Freshly ejaculated semen undergoes a process called **capacitation**: 1. inhibitory factors are washed out by uterine and fallopian fluids, 2. the sperm swims away from cholesterol vesicles, 3. the membrane of the sperms becomes more permeable to Ca^{++}


Semen

- Only 2-5 ml of semen are ejaculated, but it contains 35-200 million sperm/ml (<20 million → infertile)
- When the majority of the sperm are morphologically abnormal or nonmotile then person is likely to be infertile

Hormonal Regulation of Testicular Function


- The hypothalamus releases gonadotropin-releasing hormone (GnRH)
- GnRH stimulates the anterior pituitary to secrete FSH and LH
 - FSH causes Sertoli cells to release androgen-binding protein (ABP)
 - LH stimulates interstitial (Leydig) cells to release **testosterone**
- ABP binding of testosterone enhances spermatogenesis

HPG Axis



Hormonal Regulation of Testicular Function

- Feedback inhibition on the hypothalamus and pituitary results from:
 - Rising levels of testosterone
 - Increased inhibin



Mechanism and Effects of Testosterone Activity

- Testosterone is synthesized from cholesterol
- It binds to testosterone –binding globulin (TeBG), ABP, serum albumin, or to corticosterone-binding globulin (CBG)
- Once it diffuses to cells it either binds to androgen receptor or converted to DHT which then binds to the androgen receptor
- Testosterone targets all accessory organs, its deficiency causes these organs to atrophy
- It causes testes descent during the last 2-3 months of gestation.

Testosterone Functions:

- Testosterone targets all male reproductive organs and accessory glands, its deficiency causes these organs to atrophy
- Causes the appearance of pubic, axillary, and facial hair
- Enhances growth of the chest and deepening of the voice
- Skin thickens and becomes oily
- Bones grow and increase in density and calcium retention. It is also responsible for the male pelvis shape (narrow, long, funnel-like shape).

Testosterone functions (continued)

- It increases basal metabolic rate
- Increases red blood cells
- It also causes hair growth (pubic, axillary) and libido in females.
- Spermatogenesis and erection.

Male Sexual Act

- Erection is initiated by sexual stimuli including:
 - Touch and mechanical stimulation of the glans penis and other parts
 - Erotic sights, sounds, and smells
- Erection can be induced or inhibited solely by emotional or higher mental activity
- Enlargement and stiffening of the penis from engorgement of erectile tissue with blood

- During sexual arousal, a parasympathetic reflex promotes the release of nitric oxide, VIP, and Acetylcholine.
- Nitric oxide relaxes the penis arteries and causes erectile tissue to fill with blood
- Expansion of the corpora cavernosa:
 - Compresses their drainage veins
 - Retards blood outflow and maintains engorgement

-
- When the sexual stimulus becomes extremely intense, spinal cord begins to send sympathetic impulses to initiate emission
 - Filling of the internal urethra with semen elicits signals that promotes ejaculation
 - After orgasm, the excitement disappears within 1-2 minutes (resolution)

The Erection Reflex

